

A SIMPLE VANISHING THEOREM FOR TWISTED HOLOMORPHIC FORMS ON HERMITIAN SYMMETRIC VARIETIES

CARLA DIONISI AND DANIELE FAENZI

ABSTRACT. We find an upper bound for the degree of nonvanishing cohomology of twisted holomorphic q -forms on an irreducible Hermitian symmetric variety X as a function of q depending linearly on the twist, the dimension and the index of X , making use of simple Euclidean geometry.

1. INTRODUCTION

Let $X = G/P$ be an irreducible Hermitian symmetric variety embedded equivariantly into a projective space $\mathbb{P}(V)$ by a very ample line bundle $\mathcal{O}_X(1)$ and consider the bundle of twisted holomorphic q -forms $\Omega_X^q(\ell) = \Omega_X^q \otimes \mathcal{O}_X(1)^{\otimes \ell}$. Some vanishing theorems for the cohomology of $\Omega_X^q(\ell)$ were first obtained by Bott in [Bot57] in the case $X \simeq \mathbb{P}^n$ and extended to Grassmannians by Le Potier in [LP75] and to quadrics by Shiffman and Sommese in [SS85].

Making use of work by Kostant [Kos61], Snow [Sno86b] in [Sno86a], developed an algorithm to determine whether a given cohomology group $H^p(X, \Omega_X^q(\ell))$ vanishes for any X . These results were extended by Manivel and Snow to arbitrary homogeneous varieties in [MS96]. Applications to ample vector bundles on any projective variety can be found in [Man96], [Man97] and [Cha03].

However Snow's algorithm is of combinatoric nature and leads to nontrivial computations even on simple examples. In this paper we present a graphic and geometric approach to the problem, making use of Bott theorem and Kostant's work plus undergraduate Euclidean geometry. We find an upper bound for the degree of nonvanishing cohomology $H^p(X, \Omega_X^q(\ell))$ as a function of q, ℓ depending linearly on the dimension and the index of X .

For basic definitions and results on homogeneous varieties and representations we refer to [FH91], [Ott95], [LM03], [Tit67].

We would like to thank Giorgio Ottaviani for suggesting us these ideas.

2000 *Mathematics Subject Classification.* Primary 14M17; Secondary 14F05, 32M10, 22F30.

Key words and phrases. Homogeneous varieties;

Both authors were partially supported by Italian Ministry funds.

2. A VANISHING THEOREM FOR $H^p(X, \Omega_X^q(\ell))$

In this section we let X be an irreducible Hermitian symmetric variety i.e. a rational homogeneous variety $X = G/P$ where G is a simple algebraic group over \mathbb{C} and $P = P(\alpha_k)$ is a parabolic subgroup of G such that Ω_X^1 is an irreducible P -module. Let $\Delta = \{\alpha_1 \dots \alpha_n\}$ be the set of fundamental roots of $\mathfrak{g} = \mathfrak{g}(G)$, $\Lambda = \{\lambda_1 \dots \lambda_n\}$ be the associated set of fundamental weights, $(-, -)$ be the Killing form on \mathfrak{g} and Φ^+ be the set of positive roots of \mathfrak{g} . We have $\text{Pic}(X) \cong \mathbb{Z} = \mathbb{Z} \cdot H$ and $-K_X = \iota_X H$ where ι_X is a positive integer called the *index* of X . Define also $g = \sum \lambda_i$. Finally, for a vector μ in the weight lattice of \mathfrak{g} denote by E_μ the homogeneous bundle associated to the irreducible representation of maximal weight μ .

Theorem 2.1. *Let $X = G/P(\alpha_k)$ be an irreducible Hermitian symmetric variety and let ℓ be an integer. Define*

$$p_{\max} = \max\{p \mid \exists q \text{ such that } H^p(\Omega_X^q(\ell)) \neq 0\}$$

If $\ell \leq 0$ then $p_{\max} = 0$. If $\ell > \iota_X$ then $p_{\max} = \dim(X)$. If $0 < \ell \leq \iota_X$ then

$$p_{\max} \leq \frac{\iota_X - \ell}{2\iota_X} \dim(X)$$

3. PROOFS AND FIGURES

Lemma 3.1. *Let $X = G/P(\alpha_k)$ be an irreducible Hermitian symmetric variety. Then the expression of any weight of Ω_X^1 in the fundamental roots contains $-\alpha_k$ with multiplicity 1.*

Proof. Since T_X is the bundle associated to $\mathfrak{g}/\mathfrak{p}(\alpha_k)$, all the weights of T_X are precisely the roots of \mathfrak{g} that contain α_k with multiplicity at least 1. So any weight of Ω_X^1 contains $-\alpha_k$ with multiplicity at least 1.

Now since Ω_X^1 is indecomposable, it is given by an irreducible representation of $ss(\mathfrak{g}(\alpha_k))$, twisted by a character $\ell\lambda_k$ and $-\alpha_k$ belongs to the weight space of Ω_X^1 . Moreover the root system of $ss(\mathfrak{g}(\alpha_k))$ is generated by $\alpha_1, \dots, \hat{\alpha}_k, \dots, \alpha_n$. So any weight having a coefficient greater than 1 for $-\alpha_k$ does not lie in the $ss(\mathfrak{g}(\alpha_k))$ -orbit of $-\alpha_k$. \square

Lemma 3.2. *Let $X = G/P(\alpha_k)$ be an irreducible Hermitian symmetric variety, with $\iota_X = \iota$ and $\dim X = d$. Then the maximal weights of Ω_X^q lie*

- on the sphere $S(0)$ centered in $-g$ and passing through 0.
- on the hyperplane orthogonal to λ_k passing through the point $-\frac{q\iota}{d}\lambda_k$

Proof. Denote by H_α the hyperplane orthogonal to $\alpha \in \Phi$ and passing through $-g$ and by w_α the reflection in the weight lattice of \mathfrak{g} along the hyperplane H_α . The maximal weights of Ω_X^q are then obtained as $w_\alpha(\mu)$ where $\alpha \in \Phi^+$ contains α_k in its expression with respect to Δ and μ is a maximal weight of Ω_X^{q-1} . All such weights then lie on the sphere centered in $-g$ and since $\Omega_X^0 \cong \mathcal{O}$ the sphere passes through 0.

Now $(\alpha_i, \lambda_k) = 0$ for $i \neq k$, so the coefficient (α, μ) is a constant c for all such α 's by lemma 3.1. The weights of Ω_X^q then lie on the hyperplane orthogonal to λ_k . Notice that after d consecutive reflections 0 is mapped to $\Omega_X^d \cong K_X$ whose weight is $-\iota \lambda_k$, so the constant is $c = \iota/d$ as required. \square

Proof of theorem 2.1. Recall that $E_\mu \otimes \mathcal{O}_X(\ell) \cong E_{\mu+\ell\lambda_k}$. The maximal weights of $\Omega_X^q(\ell)$ then lie on the sphere $S(\ell)$ centered in $-g + \ell\lambda_k$ and of radius $|g|$. Consider the intersection of the two spheres $S(\ell)$ and $S(0)$. As one can read off from figure 1, or 2 these two spheres meet along the hyperplane H_z orthogonal to λ_k and passing through $z = \frac{-\iota_X + \ell}{2}\lambda_k$ i.e. the dashed line in figure 1 or 2. The figures refer respectively to the weight lattice of \mathbb{P}^2 and Q_3 .

From the proof of lemma 3.2 recall the constant $c = \iota_X / \dim(X)$. Whenever $p > \frac{\iota_X - \ell}{2c}$, the vertex of a p -th Bott Chamber μ_p (i.e. a maximal weight of Ω_X^p) lies beyond the hyperplane H_z i.e. $(\mu_p, \lambda_k) < (z, \lambda_k)$. Then, for $p > \frac{\iota_X - \ell}{2c}$ any weight of $\Omega_X^q(\ell)$ meets no Bott chamber of order p , so $H^p(\Omega_X^q(\ell)) = 0$. So the statement is proved. \square





FIGURE 1. Twisting weights of Ω^q over \mathbb{P}^2 by $\mathcal{O}(\ell)$

FIGURE 2. Twisting weights of Ω^q over Q_3 by $\mathcal{O}(\ell)$

Since irreducible Hermitian symmetric varieties are completely classified, see for instance [Ram66], one can comprehend the possible cases in the following corollary.

Corollary 3.3. *Let X be an irreducible Hermitian symmetric variety, let $0 < \ell \leq \iota_X$ be an integer and p_{\max} be defined as in theorem 2.1. Then one of the following alternatives takes place*

Group	X	$\dim(X)$	ι_X	$p_{\max} \leq$
A_n	$\mathrm{SL}(n+1)/\mathrm{P}(\alpha_{k+1})$	$(n-k)(k+1)$	$n+1$	$\frac{(n-k)(k+1)(n+1-\ell)}{2(n+1)}$
B_n	$\mathrm{Spin}(2n+1)/\mathrm{P}(\alpha_1)$	$2n-1$	$2n-1$	$\frac{2n-1-\ell}{2}$
C_n	$(2n)/\mathrm{P}(\alpha_n)$	$\frac{n(n+1)}{2}$	$n+1$	$\frac{n(n+1-\ell)}{4}$
D_n	$\mathrm{Spin}(2n)/\mathrm{P}(\alpha_1)$	$2n-2$	$2n-2$	$\frac{2n-2-\ell}{2}$
D_n	$\mathrm{Spin}(2n)/\mathrm{P}(\alpha_n)$	$\frac{n(n-1)}{2}$	$2n-2$	$\frac{n(2n-2-\ell)}{2}$
E_6	$E_6/\mathrm{P}(\alpha_6)$	16	12	$\frac{2(12-\ell)}{8}$
E_7	$E_7/\mathrm{P}(\alpha_7)$	27	18	$\frac{3(18-\ell)}{4}$

REFERENCES

- [Bot57] RAOUL BOTT, *Homogeneous vector bundles*, Ann. of Math. (2) **66** (1957), 203–248. MR 19,681d
- [Cha03] PIERRE-EMMANUEL CHAPUT, *Géométrie de quelques algèbres et théorèmes d'annulation*, Thèse de doctorat, Institut Fourier, Université Joseph-Fourier - Grenoble I., 2003.

- [FH91] WILLIAM FULTON AND JOE HARRIS, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991, A first course, Readings in Mathematics. MR **93a**:20069
- [Kos61] BERTRAM KOSTANT, *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. (2) **74** (1961), 329–387. MR 26 #265
- [LM03] JOSEPH M. LANDSBERG AND LAURENT MANIVEL, *On the projective geometry of rational homogeneous varieties*, Comment. Math. Helv. **78** (2003), no. 1, 65–100. MR **2004a**:14050
- [LP75] J. LE POTIER, *Annulation de la cohomologie à valeurs dans un fibré vectoriel holomorphe positif de rang quelconque*, Math. Ann. **218** (1975), no. 1, 35–53. MR 52 #6044
- [Man96] LAURENT MANIVEL, *Théorèmes d’annulation sur certaines variétés projectives*, Comment. Math. Helv. **71** (1996), no. 3, 402–425. MR **98a**:14030
- [Man97] ———, *Vanishing theorems for ample vector bundles*, Invent. Math. **127** (1997), no. 2, 401–416. MR **98e**:14016
- [MS96] LAURENT MANIVEL AND DENNIS M. SNOW, *A Borel-Weil theorem for holomorphic forms*, Compositio Math. **103** (1996), no. 3, 351–365. MR **97h**:32047
- [Ott95] GIORGIO OTTAVIANI, *Rational homogenous varieties*, Notes of S.M.I. course, Cortona, 1995, <http://www.math.unifi.it/ottavian>, 1995.
- [Ram66] S. RAMANAN, *Holomorphic vector bundles on homogeneous spaces*, Topology **5** (1966), 159–177. MR 32 #8357
- [Sno86a] DENNIS M. SNOW, *Cohomology of twisted holomorphic forms on compact Hermitian symmetric spaces*, Représentations des groupes et analyse complexe (Luminy, 1986), Journées SMF, vol. 24, Univ. Poitiers, Poitiers, 1986, pp. 77–87. MR **89b**:32042
- [Sno86b] ———, *Cohomology of twisted holomorphic forms on Grassmann manifolds and quadric hypersurfaces*, Math. Ann. **276** (1986), no. 1, 159–176. MR **87m**:32066
- [SS85] BERNARD SHIFFMAN AND ANDREW JOHN SOMMESE, *Vanishing theorems on complex manifolds*, Progress in Mathematics, vol. 56, Birkhäuser Boston Inc., Boston, MA, 1985. MR **86h**:32048
- [Tit67] JACQUES TITS, *Tabellen zu den einfachen Lie Gruppen und ihren Darstellungen*, Springer-Verlag, Berlin, 1967. MR 36 #1575

DIPARTIMENTO DI MATEMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A,
I-50134, FLORENCE, ITALY

E-mail address: dionisi@math.unifi.it

URL: <http://www.math.unifi.it/~dionisi/>

DIPARTIMENTO DI MATEMATICA “U. DINI”, UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A,
I-50134, FLORENCE, ITALY

E-mail address: faenzi@math.unifi.it

URL: <http://www.math.unifi.it/~faenzi/>